

Self-stabilization of Circular Arrays of Automata*

Leonid A. Levin[†]
Boston University[‡]

Keywords: self-stabilization, error-correction, cellular automata.

As computing elements shrink closer to molecular size, the reliability becomes a serious issue. John von Neumann proposed circuits which compute reliably in the presence of noise. They cannot be realized, however, with local connections on an Euclidean grid. Andrei Toom designed a planar grid of simple automata which stores one bit reliably despite noise. Aligned in layers they were used by Peter Gacs and John Reif to simulate a Turing Machine on a three-dimensional grid. The need to dispose of heat restricts computing chips to 2 dimensions, which would make this method to depend on linear arrays of automata to preserve each bit reliably. Such automata were believed impossible until Kurdiumov and Gacs constructed them in a series of monumental works. Their complexity is enormous, which leaves a need for simpler solutions.

[Gacs, Kurdiumov, Levin, 78] proposed simple automata with 2 states “ $\triangleright, \triangleleft$ ”. They are arranged in a linear array and change state when opposed by both first and third neighbor from the sharp end. In an infinite array they are self-stabilizing: if all but a finite minority of automata are in the same state, the minority states disappear. Implicit in the paper was a stronger result that a sufficiently small minority of states vanish even in a finite circular array (ring). I think this strengthening deserves to be made explicit which is the purpose of the following note.

Theorem 1 . *Let $\alpha = 1/x \approx 1.7734$ where $5^x = 2^x + 1$. Let in a ring of more than $3k^\alpha$ GKL automata all but k of them start in the same state. Then the minority states disappear within $3k^\alpha$ steps.*

Assume, the $\triangleright$ -s are in majority. We refer as intervals to maximal strings of three or more $\triangleright$ -s and as segments to strings with $\triangleleft$ at each end. Let S be a segment between $\triangleright^{2|S|+1}$ and $\triangleright^{|S|+2}$. It grows at most one cell per step to the left and never to the right. The left end of its leftmost pattern $\triangleleft\triangleleft$ or $\triangleleft\triangleright\triangleright\triangleleft$ moves right each step until gone. Then S shrinks at the right end by at least 3 cells per step. So, $2|S|$ steps erase any effect of S . We say S is *killed* by the surrounding $\triangleright$ intervals. Any segment S is killed unless it has a $\triangleleft$ within either $|S| + 1$ cells to the right or $2|S|$ to the left. We say S is closed to this side and break a ring into a binary hierarchy of *solid* segments, each of which can be killed only as a whole: We start with single $\triangleleft$ -s and combine into a higher solid any two solid neighbors closed toward each other.

Let S be a counterexample to the Theorem with the fewest $\triangleleft$ -s. Its maximal solids must all be closed from one side (and so occupy at least a third of S). This side must be left: If all are open to the left, then the left neighbor of the longest interval is killed; if some are closed to the left and some to the right, they combine further. Take the solid P next right to the longest on S interval p . Expanding left, P sheds its $\triangleleft\triangleleft$ and $\triangleleft\triangleright\triangleright\triangleleft$ patterns and shrinks moving left until it hits its left neighbor and is overrun. Its right end moves left for at least $2|p| - |P| - 2$ cells, extending the original $\geq |P| + 2$ cells of the interval at its right enough to kill its right neighbor. Thus, S can have only one maximal solid, closed on both sides, and $|S| \leq 2|P| + 1$.

It is left to prove that no solid with k $\triangleleft$ -s has more than $(3k^\alpha - 1)/2$ cells. Consider a shortest solid which violates that. It consists of two solids of n and $nt - 1 \geq n$ cells respectively and an interval of $nr \leq n \min\{2, t\}$ cells. It has at least $((2n + 1)/3)^x + ((2nt - 1)/3)^x \geq (2n/3)^x(1 + t^x)$ $\triangleleft$ -s. We must see that this is at least $((2n + 2nr + 2nt - 1)/3)^x < (2n/3)^x(1 + r + t)^x$, i.e. that $1 + t^x - (1 + t + \min\{2, t\})^x \geq 0$. Since $t > 1$, this function has its minimum 0 at $t = 2$. ■

The factor 3 can be improved and David Metcalf conjectured that the power can be improved too, to the $1/\log$ of the golden ratio ≈ 1.44 . It certainly cannot be improved beyond that: The strings $S_0 = S_1 = \triangleleft^3$, $S_{i+2} = S_i \triangleright^{|S_i|-4} S_{i+1}$ with $|S_i| = 2^{i-1} + 2$ have 3 Fibonacci(i) $\triangleleft$ -s and kill all their $\triangleright$ -s.

* *Theor. Comp. Sci.*, 235(1):143-144, 3/17/2000.

[†]Supported by NSF grants CCR-9610455, CCR-9820934.

[‡]Computer Science department, 111 Cummington St., Boston, MA 02215; (e-mail to Lnd@bu.edu)

References

- [1] Peter Gács. Reliable computation with cellular automata. *J. Comp. System Sci.* 32(1):15-78, 1986.
- [2] P. Gács, G.L. Kurdyumov, L. Levin. One-Dimensional Homogeneous Media Dissolving Finite Islands. *Problems of Information Transmission* (trans. from *Problemy Peredachi Informatsii* (in Russian)), 14(3):223-226, 1978.
- [3] G.L. Kurdyumov. An Example of a Nonergodic Homogeneous One-Dimensional Random Medium with Positive Transition Probabilities. *Soviet Math. Doklady* 19(1):211–214, 1978.
- [4] John von Neumann. Probabilistic logics and the synthesis of reliable organisms from unreliable components. In C. Shannon, McCarthy, eds., *Automata Studies*, Princeton University Press, Princeton NJ, 1956.
- [5] Andrei Toom. Monotonic binary cellular automata. *Problems of Information Transmission* (trans. from *Problemy Peredachi Informatsii* (in Russian)), v. 12, 1976.